

Physiology

KEYWORDS: metabolic syndrome, exercise, physical activity

STUDY OF PHYSICAL EXERCISE AND METABOLIC SYNDROME



Volume - 6, Issue - 9, September - 2021

ISSN (O): 2618-0774 | ISSN (P): 2618-0766

Dr. Ramanuj

Junior Resident, Deptt of pulmonary medicine PMCH Patna.

Dr. Divya Gupta*

Junior Resident, Deptt of physiology AIIMS Rishikesh. *Corresponding Author

INTERNATIONAL JOURNAL
OF PURE MEDICAL RESEARCH

**ABSTRACT**

Regular physical activity practice has been recommended for the prevention and rehabilitation of cardiovascular diseases and other chronic diseases by different health care associations world-wide. The regular practice of physical exercise has been shown to have beneficial effects in the prevention and treatment of blood hypertension, insulin resistance, diabetes, dyslipidemia, and obesity. Physical training therefore should be encouraged for both healthy individuals and those with multiple risk factors if they are capable of participating in a physical fitness program. Just as clinical therapy helps to maintain the function of organs, physical activity promotes favorable physiological adaptations that result in an improved quality of life.

Physical inactivity and low physical conditioning level have been considered as risk factors for early mortality so important as the smoking, dyslipidemia and arterial hypertension. Epidemiological studies have demonstrated direct relation between physical inactivity and the presence of cardiovascular risk factors such as arterial hypertension, insulin resistance, diabetes, dyslipidemia and obesity. On the other hand, the regular practice of physical activity has been recommended for the prevention and treatment of cardiovascular disease, their risk factors and other chronic diseases.

The metabolic syndrome – also known as syndrome X, insulin resistance syndrome, deadly quartet or plurimetabolic syndrome – is characterized by the group of cardiovascular risk factors such as arterial hypertension, insulin resistance, hyperinsulinaemia, glucose intolerance/diabetes type II, central obesity and dyslipidemia (high LDL-cholesterol, high triglycerides and low HDL-cholesterol). Epidemiological and clinical studies have demonstrated that the regular practice of physical activity is an important factor for prevention and treatment of this disease.

The objective of this reviewing is to demonstrate the role of the regular practice of physical activity in the prevention and treatment of the metabolic syndrome as well as to describe the amount and modality of exercise required for this purpose.

EXERCISE AND OBESITY

A rapid and increasing growth of obese people has been observed, what made the obesity a public health problem. This disease has been classified as a disorder primarily of high energetic ingestion. However, there are evidences suggesting that most cases of obesity are more related to the low energy expenditure than to the high food ingestion, where the physical inactivity of the modern life seems to be the highest etiological factor for the growth of this disease in industrialized societies.

Epidemiological and cohort studies have demonstrated strong association between obesity and physical inactivity. Inverse as-

sociation between physical activity, body mass index (BMI)*, hip-waist ratio (HWR)* and waist circumference has also been reported. These studies demonstrate that the benefits of physical activity on obesity may be reached with low, moderate or high intensities, indicating that the maintenance of an active lifestyle, regardless of what physical activity is performed, may avoid the development of this disease.

It is required that the energy expenditure be higher than the daily energy intake, what leads us to think that a simple reduction on the amount of food through an alimentary diet is sufficient. However, it is not so simple and studies demonstrate that the change in the lifestyle through the increases on the amount of physical activity practiced and alimentary reeducation is the best treatment.

The energetic expenditure is composed of three main components: rest metabolic rate (RMR), thermic effect of physical activity and thermic effect of food (TEF). The RMR, which is the energetic cost to maintain systems functioning in rest, is the component of higher daily energetic expenditure (60 to 80% of the total). The obesity treatment only through dietary caloric restriction leads to a decrease on the RMR (through the decrease on the muscular mass) and on the TEF, what leads to a reduction or maintenance of weight reduction and to a tendency of returning to initial weight, despite the continuous caloric restriction, thus contributing to a long-lasting poor efficiency of this intervention. However, the combination of caloric restriction and physical exercise helps to maintain RMR, improving the results of long-term weight reduction programs.

This occurs because physical exercise elevates RMR after its performance due to the increase on the substrate oxidation, catecholamine level and stimulation of protein synthesis. This effect of the exercise on RMR may last from 3 hours until 3 days, depending on the type, intensity and duration of the exercise.

Activity	45 kg	68 kg	90 kg
Pedaling 10 km/h	160	240	312
Walking 3.2 km/h	160	240	312
Walking 4.8 km/h	210	320	416
Walking 7.2 km/h	295	440	572
Trotting 11 km/h	610	920	1,230
Running 16 km/h	850	1,280	1,660
Swimming	185	275	385

Figure 1 – Approximate energetic expenditure per hour of a person (45, 68 and 90 kg) performing physical activity

Although most studies have investigated the effect of the physical exercise on the weight reduction, the inclusion of the weight-resistance exercise (weightlifting) shows many advantages. The weight-resistance exercise is a powerful stimulus to increase mass, strength and muscular power also helping to preserve musculature, which trends to decrease due to diet, maximizing the reduction of the body fat. Furthermore, its potential in improving strength and muscular resistance may be especially positive in daily tasks, furthering the adoption of a more active lifestyle in inactive obese individuals.

The traditional recommendation of at least 150 weekly minutes (30 minutes, 5 times a week) of physical activity from light to moderate intensities, which is primarily based on the effects of the physical activity on the cardiovascular disease and other chronic disease such as diabetes, demonstrates not being sufficient for weight reduction programs. Thus, it has been recommended that exercise programs for obese individuals start with a minimum of 150 weekly minutes in moderate intensity progressively increasing up to 200 to 300 weekly minutes at the same intensity. However, if for any reason the obese could not reach this target, he should be encouraged to perform at least the minimum recommendation of 150 weekly minutes, once with no weight reduction, there will be benefits to health.

the increased and continuous utilization of fatty acids as muscular fuel. However, these are extreme conditions in which the exercise intensity is higher than the intensity that most individuals with metabolic syndrome can stand.

The effect of the physical exercise on the insulin sensibility has been demonstrated from 12 to 48 hours after the exercise session, however, it returns to pre-activity levels in 3 to 5 days after the last physical exercise session, what reinforces the necessity to practice physical exercises with frequency and regularity.

The fact that only one exercise session improves the insulin sensibility and that the effect provided by the training withdraws in a few days of inactivity arises the hypothesis that the physical exercise effect on the insulin sensibility is merely acute. However, it has been demonstrated in a study in which individuals with insulin resistance improved the insulin sensibility in 22% after the first exercise session and in 42% after 6 weeks of training, what demonstrates that the physical exercise presents both an acute effect and a chronic effect on the insulin sensibility.

Exercise and diabetes type 2

Epidemiological and intervention studies clearly demonstrate that the regular practice of physical exercises is effective for the prevention and control of diabetes type 2. The regular practice of physical exercises have demonstrated to decrease the risk of developing diabetes type 2 both in men and women, regardless the family history, weight and other cardiovascular risk factors such as smoking and hypertension.

Intervention studies have demonstrated that changes in the lifestyle with the adoption of new alimentary habits and the regular practice of physical exercises decrease the incidence of diabetes type 2 in individuals with glucose intolerance, where the performance of at least four weekly hours of physical activity from moderate to intense intensity decreased 70% the incidence of diabetes type 2 on average, in relation to the inactive lifestyle after four years of follow-up.

The practice of physical activity has also been considered as an important tool in the treatment of individuals with diabetes type 2. Physical exercise programs have demonstrated to be effective in the glycemic control of diabetic individuals, improving the insulin sensibility and the glucose tolerance and decreasing the blood glycemiy of these individuals.

The performance of aerobic exercises have been generally recommended for individuals with diabetes type 2. However, recent studies have demonstrated that the weight-resistance exercise is also beneficial in the glycemic control of individuals with diabetes type 2.

The weight-resistance exercise may be especially useful for aged diabetic individuals, once during aging, the muscular strength and mass decreases, undesirably affecting the energetic metabolism (figure 2). The increase on the muscular mass and strength through the practice of weight-resistance exercises may revert this situa-

tion, improving the glycemic control of these individuals. Thus, decreases on the blood glucose levels, increases on the muscular glycogen supply, reduction on the systolic pressure and trunk fat have been demonstrated as well as increases on the muscular mass and daily physical activity levels of aged diabetic individuals from both gender after 16 weeks of weight-resistance exercises, what resulted in reduction on dedication up to 72%, while individuals who participated in the control group presented unchanged blood glycemiy levels, systolic pressure, trunk fat and daily physical activity and decreased muscular glycogen supplies and 42% had increased medication.




Figure 2 – Effect of the strength and muscular mass reduction occurring with aging in several metabolic variables associated to metabolic syndrome. Adapted from Ciolac and Guimarães with the authors' consent.

EXERCISE AND ARTERIAL HYPERTENSION

Epidemiological and clinical studies have demonstrated beneficial effects of the practice of physical exercises on the arterial pressure in individuals of all ages. High level of daily physical activity is associated to lower levels of arterial pressure in rest. The regular practice of physical exercises have demonstrated to prevent blood pressure increases associated to age even in individuals with increased risk to develop it. Physical activity programs have demonstrated to decrease the systolic and diastolic blood pressure both in hypertensive and normotensive individuals.

These benefits of the physical activity on blood pressure make physical activity an important tool on prevention and treatment of the hypertension. A meta-analysis of 54 controlled randomized longitudinal studies that investigated the effect of the aerobic physical activity on the blood pressure demonstrated that this exercise modality reduces 3,8 mmHg and 2,6 mmHg the systolic and diastolic pressures on average, respectively. Reductions of only 2 mmHg on the diastolic pressure may reduce significantly the risk of diseases and deaths associated to hypertension, what demonstrates that the practice of aerobic exercise represents important benefit for health in hypertensive individuals.

It has been proposed that the effect of the aerobic exercise on the blood pressure is more due to the acute effect of the last exercise session than to the training cardiovascular adaptations. In this context, the study performed by our research group demonstrated that hypertensive individuals presented reductions on the blood pressure clinical monitoring during 24 hours (MAPA 24 h) when performed shortly after the last exercise session, which were not observed when performed 72 hours after the last session.

Hypertensive individuals have been traditionally discouraged to perform weight-resistance exercises due to the fear that this exercise modality would cause a cerebrovascular or cardiac event.

However, studies that investigated the effect of long periods of weight-resistance exercise training on the blood pressure in rest reported no harmful effects, suggesting that hypertensive individuals should not avoid its practice, once it provides relevant benefits to quality of life, especially for aged individuals.

EXERCISE AND DYSLIPIDEMIA

The effects of the physical activity on the lipids and lipoprotein profiles are well known. Individuals physically active present higher levels of HDL-cholesterol and lower levels of triglycerides, LDL (65) and VLDL-cholesterol, if compared to inactive individuals. Intervention studies demonstrate the unfavorable lipids and lipoprotein profiles improve with physical training. These improvements are not dependent on gender, body weight and diet, however, there is a possibility of being dependent on the glucose tolerance degree. The physical activity has demonstrated to be effective in decreasing the level of VLDL-cholesterol in individuals with diabetes type 2, however, except for a few, most studies have not demonstrated significant improvement on levels of HDL and LDL-cholesterol in this population maybe due to the low intensity of the exercise employed.

Although studies on the effect of physical exercises on the lipids and lipoprotein profiles in individuals with metabolic syndrome are scarce, considering the evidences above and the fact that physical exercises increase the ability of the muscular tissue in spending fatty acids and the activity of the enzyme lipoprotein lipase in the muscle, it is likely that the physical exercise be effective in improving the lipid and lipoprotein profiles in individuals with metabolic syndrome.

PRESCRIPTION OF PHYSICAL EXERCISE

In the practice or prescription of physical training with the objective of obtaining some training physiological effect, either the improvement of the physical conditioning or the prevention and treatment of diseases, one should take into account four basic principles. The first is the overload principle that commends that in order to obtain physiological response with physical training, the physical exercise should be performed with an overload above the usual, which could be controlled with intensity, duration and exercise frequency. The second principle is the specificity principle, which is characterized by the fact that exercise specific modalities unchain specific adaptations that further specific physiological responses. The third principle is the individuality principle, where the biological individuality of each group should be respected in the prescription of some exercise program, once the same overload and exercise modality will cause responses of different magnitude in different individuals. The fourth and last principle is the reversibility principle, which is characterized by the fact that the physiological adaptations furthered by the physical exercise performance return to the pre-training original state, when the individual returns to the inactive lifestyle.

The performance of at least 30 minutes of physical activity (performed formally or in the spare time, continuously or concentrated in sessions of at least 10 minutes) of intensity at least moderate (level 12 in the Borg scale – fig. 3) performed in most days of the

6.
7. Very easy
8.
9. Easy
10.
11. Relatively easy
12.
13. Slightly tiring
14.
15. Tiring
16.
17. Very tiring
18.
19. Exhaustive
20.

Figure 3 – Borg scale of effort subjective perception

in order for the benefits and health security of the regular practice of physical activity to be maximized, a prescription of exercises that takes into account the necessities, aims, initial capacities and family history is required. Furthermore, for the treatment of some diseases such as obesity, this amount of exercises has demonstrated being not sufficient. These factors lead us to believe that the individual with metabolic syndrome, since they present risk factor for cardiovascular diseases, will obtain higher benefits with the regular practice of physical exercise if it is individually planned, aimed at the improvement of his health state and taking into account his health state, risk factors and physical capacity as well as his history and aims.

The amount of researches on the effects of the weight-resistance training on metabolism is small if compared to researches involving aerobic exercises.

However, based on current reviewing studies we propose that a physical activity program aimed at the prevention and treatment of the metabolic syndrome should include components that improve the cardiorespiratory conditioning, strength and muscular resistance.

– Both weight-resistance and aerobic exercises further substantial benefits related to health and physical conditioning (figure 4) including most risk factors of the metabolic syndrome.

Variable	Aerobic exercise	W. resistance exercise
Glucose metabolism		
Glucose tolerance	↓↓	↓↓
Insulin sensibility	↑↑	↑↑
Serid lipids		
HDL cholesterol	↑ ↔	↑ ↔
LDL cholesterol	↓ ↔	↓ ↔
Blood pressure in rest		
Systolic	↓ ↔	↔
Diastolic	↓ ↔	↓ ↔
Body composition		
% of fat	↓↓	↓
Body mass free of fat	↔	↑↑
Basal metabolism	↑	↑↑↑
Muscular strength	↔	↑↑↑
Aerobic capacity		
vo2max	↑↑↑	↑ ↔
Time of maximal or submaximal aerobic exercise	↑↑↑	↑↑
↑ = increase on values; ↓ = reduction on values; ↔ = unchanged values; ↑ or ↓ = small effect; ↑↑ or ↓↓ = intermediate effect; ↑↑↑ or ↓↓↓ = large effect; HDL cholesterol = high-density cholesterol; LDL cholesterol = low-density cholesterol.		

Figure 4 – Effect of the aerobic and weight-resistance exercise on variables that influence the metabolic syndrome and physical conditioning

The mechanisms through which weight-resistance and aerobic exercises affect some variables of the metabolic syndrome such as the insulin resistance, the glucose intolerance and obesity seem to be different and the sum of the effects from both activities may occur.

Although the minimal dose of exercise required to reach many of

the health benefits is known, the optimum dose for prevention and treatment of most disorders is yet unknown.

With regard to aerobic exercises, it has been recommended to be performed from 3 to 6 times a week with intensity from 40 to 85% FCR (40 to 85% of the VO₂ or 55 to 90% of the FC_{max} or level 12 to 16 in the Borg scale) and duration of 20 to 60 minutes. Due to the fact that higher exercise intensities is associated to higher cardiovascular risk and orthopedic lesions and the lower adherence to physical exercise programs, it is recommended that programs aimed at inactive individuals with risk factor for cardiovascular diseases emphasize moderate intensity (50 to 70% of the FCR and levels 12 to 13 in the Borg scale) and long duration (30 to 60 minutes)

The current recommendation for the practice of weight-resistance exercise is 1 series of 8 to 12 repetitions (10 to 15 for individuals above 50/60 years of age), intensity of 50 to 70% of the maximal load (13 to 15 in the Borg scale) performed with 8 to 10 exercises working all large muscular groups 2 to 3 times a week. However, this recommendation is only based on strength improvement and muscular resistance. The performance of a higher number of series (2 to 3) will increase the energetic expenditure of the exercise session, increasing the activity benefit for individuals with metabolic syndrome.

Thus, we recommend that individuals with metabolic syndrome start with 1 series, progressively increasing up to 2 and 3 series after adaptation.

All aerobic and weight-resistance physical activity sessions should include warm-up and calm down with the use of flexibility exercises at the beginning and at the end of each session.

REFERENCES

- Blair SN, Kampert JB, Kohl III HW, Barlow CE, Macera CA, Paffenbarger RS, et al. Influences of cardiorespiratory fitness and other precursors on cardiovascular disease and all-cause mortality in men and women. 1996;276:205-10.
- Rennie KL, McCarthy N, Yazdgerdi S, Marmot M, Brunner E. Association of metabolic syndrome with both vigorous and moderate physical activity. 2003;32:600-6.
- Gustat J, Srinivasan SR, Elkasabany A, Berenson GS. Relation of self-rated measures of physical activity to multiple risk factors of insulin resistance syndrome in young adults: Clin Epidemiol 2002;55:997-1006.
- Wareham NJ, Hennings SJ, Byrne CD. A quantitative analysis of the relationship between habitual energy expenditure, fitness and the metabolic cardiovascular syndrome. Br J Nutr 1998;80:235-41.
- Lakka TA, Laaksonen DE, Laaka HM, Männikö N, Niskanen LK, Raumamaa R, et al. Sedentary life style, poor cardiorespiratory fitness, and the metabolic syndrome. Med Sci Sports Exerc 2003;35:1279-86.
- Pate RR, Pratt M, Blair SN, Haskell WL, Macera CA, Bouchard C, et al. Physical activity and public health: a recommendation from the Centers for Disease Control and Prevention and the American College of Sports Medicine. JAMA 1995; 273:402-7.
- Fletcher GF, Balady GJ, Amsterdam EA, Chaitman B, Eckel R, Fleg G, et al. Exercise standards for testing and training: a statement for healthcare professionals from the American Heart Association. Circulation 2001;104:1694-1740.
- NIH Consensus Development Panel on Physical Activity and Cardiovascular Health. Physical activity and cardiovascular health (NIH consensus conference). JAMA 1996;276:241-6.
- Department of Health and Human Services. Physical activity and health: a report of Surgeon General. Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, 1996.
- Sociedade Brasileira de Cardiologia. III Diretrizes brasileiras sobre dislipidemias e diretriz sobre prevenção da aterosclerose do Departamento de Aterosclerose da Sociedade Brasileira de Cardiologia. Bras Cardiol 2001:77-98.
- Paffenbarger RS, Jung DL, Leung RW, Hude RT. Physical activity and hypertension: an epidemiological view. Ann Med 1991;23:319-27.
- Whelton SP, Chin A, Xin X, He J. Effect of aerobic exercise on blood pressure: a meta-analysis of randomized, controlled trials. Ann Intern Med 2002;136:493-503.
- Eriksson J, Taimela S, Koivisto VA. Exercise and the metabolic syndrome. Diabetologia 1997;40:125-35.
- Castaneda C, Layne LE, Orians LM, Gordon PL, Walsmith J, Foldvari M, et al. A randomized controlled trial of resistance exercise training to improve glycemic control in older adults with type 2 diabetes. Diabetes Care 2002;25:2335-41.
- Durstine JL, Haskell WL. The influence of exercise on plasma lipids and lipoproteins. Exerc Sport Sci Rev 1994;22:477-521.
- American College of Sports Medicine. ACSM stand position on the appropriate intervention strategies for weight loss and prevention of weight regain for adults. Med Sci Sports Exerc 2001;33:2145-56.
- Bielinski R, Schutz Y, Jéquier E. Energy metabolism during the post-exercise recovery period in man. Am J Clin Nutr 1985;42:69-82.
- Horton ES. Metabolic aspects of exercise and weight reduction. Med Sci Sports Exerc 1985;18:10-8.
- Tremblay A, Nadeau A, Fournier G, Bouchard C. Effect of a three-day interruption of exercise training on resting metabolic rate and glucose-induced thermogenesis in trained individuals. Obes 1988;12:163-8.
- McArdle WD, Katch FI, Katch VL. Fisiologia do exercício: energia, nutrição e desempenho humano. 4ª ed. Rio de Janeiro: Ed. 1998.
- Baalor DL, Katch VL, Becque MD, Marks CR. Resistance weight training during caloric restriction enhances lean body weight maintenance. Nutr 1988;47:19-25.
- Geliebter A, Maher MM, Gerace L, Gutin B, Heymsfield SB, Hashim SA. Effects of strength or aerobic training on body composition, resting metabolic rate, and peak oxygen consumption in obese dieting subjects. Nutr 1997;66:557-63.