

Cardiology

KEYWORDS:

ano,High,Low,Seton,Fistulectomy,
Fistulotomy,Dressing,Antibiotics

STUDY OF MANAGEMENT OF 25 CASES OF FISTULA IN ANO



Volume - 5, Issue - 2, February - 2020

ISSN (O): 2618-0774 | ISSN (P): 2618-0766

Dr Vikas Warikoo

(MS,MCh)

Dr Sandipkumar
Chaudhari*

(MS,FMAS,FIAGES.FCCS) *Corresponding Author drsandy253@gmail.com

Dr Adwait Patel

(MBBS)

INTERNATIONAL JOURNAL
OF PURE MEDICAL RESEARCH

**Abstract:****Introduction:-**

Fistula-in-ano forms a good majority of treatable benign lesions of the rectum and anal canal. 90% or so of these cases are end results of crypto

glandular infections. These conditions affect a lot of young and middle aged persons causing loss of valuable productive man hours.

Aims:

To know the usefulness of investigative procedures in early and accurate diagnosis of fistula in ano. To study the efficacy of different modalities of surgical approach with reference to post-operative hospital stay and complications like pain, bleeding and sphincteric incontinence and outcome in respect to persistence/ recurrence of fistulae.

Materials & Methods:-

Cross sectional were observational study of 25 patients with fistulae in Ano admitted Surgical ward of a tertiary care

hospital of age between 20 years and 80 years of age.

Conclusion :-

Fistula in ano is an important, commonest disease due to crypto glandular infection (anal glands) and has a complication of ano rectal abscess. It is curable disease by the treatment of surgery and higher antibiotics, local antibiotics with good post operative wound management, like sitz bath for twice a day without closing the wound.

INTRODUCTION

Fistula-in-ano forms a good majority of treatable benign lesions of the rectum and anal canal. 90% or so of these cases are end results of crypto glandular infections. As such, the vast majority of these infections are acute but a significant minority is contributed by chronic, low-grade infections, hence pointing to varying etiologies. The common pathogenesis however is the bursting open of an acute or inadequately treated ano-rectal abscess into the peri-anal skin. Most of these fistulas are easy to diagnose with a good

source of light, a proctoscope, and a meticulous digital rectal examination. Despite the easy of diagnosis, establishing a cure is problematic on two accounts. Firstly, many patients tend to let their ailment nag them rather than being subject to examination, mostly owing to the site of this disease. The more important second factor is

that a significant percent of these diseases persist or recur when the right modality of surgery is not adopted or when the post-operative care is inadequate. So these conditions affect a lot of young and middle aged persons causing loss of valuable productive man hours.

MATERIAL AND METHODS

A total of 25 patients with clinically diagnosed fistula in ano were included in the study. All patients were subjected to surgical intervention. The study was conducted at tertiary care Hospital during period of 2009 to 2012. Clinical history was obtained in all the patients. Clinical examination including per rectal examination and proctoscopy was done in all the patients. All the patients were processed by routine investigations, ECG, chest X - ray etc prior to surgery. Patients were followed up to a period of 1 year.

Cases were selected by following criteria:

Inclusion criteria:

- 1) Age at time of admission between 20yrs and 80yrs
- 2) Patient complaint of recurrent discharge per rectum for a period greater than 15 days with demonstration of single/multiple external openings seen as an elevation of granulation tissue discharging pus in the perianal region and confirmed on digital rectal examination.

Exclusion criteria:

- 1) Patients with clinical and investigative evidence of ano-rectal malignancy.
- 2) Patients refusal for surgical intervention when the fistula in ano was demonstrated on clinical examination.

Level of fistula

The internal opening was demonstrated by digital rectal examination, proctoscopy, injection of hydrogen peroxide or dilute methylene blue through the external opening or fistulogram.

- 1) Low level fistula had internal opening situated below the ano rectal ring.
- 2) High level fistula had internal opening situated above the ano rectal ring.

Fistulogram was performed in the following cases:

- 1) Recurrent fistulas
- 2) Patients with clinical and investigative evidence of tuberculosis/inflammatory bowel disease
- 3) Demonstration of multiple external openings on inspection of perianal region and confirmed on digital rectal examination.
- 4) The internal opening was not apparent on digital rectal examination and proctoscopy.

TREATMENT

- 1) Patients with low level fistula were treated with fistulotomy and fistulectomy.

2) Patients with high level fistula were treated with seton placement.

POST OPERATIVE OUTCOME:

Different modalities of surgical approach were studied with reference to:

- 1) Duration of hospital stay
- 2) Complications
- a) Bleeding
- b) Pain
- c) Urinary retention
- d) Incontinence
- e) Recurrence

OBSERVATION & DISCUSSION

25 cases of fistula in ano were selected randomly and studied in detail the following results were obtained.

1.Age incidence

Age	20-25	26-30	31-35	36-40	41-45	46-50	51-55	>55
No.	6	6	2	3	3	4	1	0
%	24	24	8	12	12	16	4	0

Only patients with age at time of admission between 20yrs and 80yrs were included in the study as other age groups were less convinced for surgery. In our study fistula in ano was more common(48%) in 20-30 years age group.

Kyung Won Kang, M.D., Kyung Lim Choi, and Hong Kyun Kim, in their study “A Clinical Study of Fistula-in-ano” had majority of patients in the third to fifth decade of life (75.1 %).

2. Previous history for fistula in ano

6 patients i.e. 24% had similar illness out of them two previously operated for fistula with recurrence, and four patients with similar illness and resolved without treatment.

Inadequately operated patient may show healing of wound from the superficial side of track but rectal and anal side of wound may persist leading to recurrence, so healing from depth is very necessary.

3. No. of external openings of Fistula in ano


No. of external openings	No. of patients	%
Single	22	88
Multiple	3	12

In present study, 88% of patients had only one external opening and 12% of patients had multiple external openings.

Kyung Won Kang, M.D., Kyung Lim Choi, and Hong Kyun Kim, in their study “A Clinical Study of Fistula-in-ano” demonstrated that the majority of patients had one external opening(88.0%).

From this fact we can conclude that fistula in ano is present in majority of cases, with only one external opening

4. Fistulogram



Fistulogram was done in only 10 cases i.e. 40% of cases in this series. This was done where clinical impression of type of fistula could not

be made confidently or fistula was associated with complicating factors. Fistulogram has high degree of inaccuracy and is invasive and potentially may result in the dissemination of sepsis, so it should be used judiciously in evaluation of fistula.

5.Level of fistula


Type of fistula	Cases	%
Low anal	18	72
High anal	7	28

In this study, 72% of patients had low level of fistula and another 28% of patients had an internal opening situated above the ano rectal ring. Patients with low level fistula were treated with fistulotomy and fistulectomy and patients with high level fistula were treated with seton placement.

6.Types of Surgical Treatment

Type of operation	Cases	%
Fistulotomy	15	60
Fistulectomy	3	12
Seton placement	7	28

In this study 60 % of patients underwent Fistulotomy, 12 % of patients Fistulectomy and another 28% seton placement. Patients with low level fistula were treated with fistulotomy and fistulectomy and patients with high level fistula were treated with seton placement.



G. Rosa¹, P. Lolli¹, D. Piccinelli¹, F. Mazzola¹ and S. Bonomo¹ in their study “Fistula in ano: anatomoclinical aspects, surgical therapy and results in 844 patients” had treated most patients by fistulotomy alone (594 patients, 70.4%) or by the combined fistulectomy-fistulotomy method (237 patients, 28.1%), with or without loose seton.

Kim JW, Kwon SW, Son SW, Ahn DH, Lee KP study of “Comparative Review of Perianal Sinus & Fistula in Ano” had used fistulectomy in 4 cases (23.5%), fistulotomy in 11 cases (64.7%) and seton procedure in 2 cases (11.8%).

Fistulectomy creates larger wounds significantly prolonging wound healing time and there is greater risk of injuring or excising underlying muscle thereby increasing the risk of incontinence. So fistulectomy was preferred in fewer number of

patients. Seton when tightened cuts through the anal sphincter musculature gradually and also the tract cut under goes fibrosis with resultant healing of the fistula. As a result there is less chances of incontinence and hence was used in our study for management of high fistula.


7. Average duration of post operative hospital stay(in days)

Type of operation	Average duration of post operative hospital stay(in days)
Fistulotomy	5
Fistulectomy	11
Seton placement	12

Seton placement was associated with the maximum duration of post operative hospital stay(average 12 days) followed by fistulectomy with 11 days. Fistulotomy required the least days of post operative hospital stay with an average of 5 days.

8. Post Operative Complications

Type of operation	Post Operative Complications rate (%)				
	Bleeding	Pain	Urinary retention	Incontinence	Recurrence
Fistulotomy	7	7	7	nil	7
Fistulectomy	33	67	nil	33	nil
Seton placement	nil	29	nil	nil	nil



a) Bleeding

Post operative bleeding occurred in 7% of fistulotomy patients and 33% in the fistulectomy group. No clinical significant bleeding occurred in the seton group. Bleeding was controlled in all the cases with pressure dressing only and no further operative management was required.

b) Pain

7% of fistulotomy patients ,67% of fistulectomyand 29% of seton group complained of severe pain. Pain incidence was significantly higher in fistulectomy and seton group as compared to fistulotomy group. Pain was treated by analgesics and local anaesthetic ointment application.

c) Urinary retention

Retention of urine was noticed in 7% of fistulotomy patients. No similar complaints were noticed in other groups. Retention was relieved by indwelling catheterisation.

d) Incontinence

Incontinence of flatus was noticed in 33% of fistulectomy patients and was absent in the other groups. None of the patients in our study developed incontinence of stools.

Wolfram Trudo Knoefel, Stefan B. Hosch, Björn Hoyer, Jakob R. Izbicki study “The Initial Approach to Anorectal Abscesses: Fistulotomy Is Safe and Reduces the Chance of Incontinence” had 4 of the 131 patients (3%) developing incontinence of liquid stool and flatus, but no incontinence of solid stool occurred in their study as well.

R. D. Ritchie, J. M. Sackier and J. P. Hodde in their study “Incontinence rates after cutting seton treatment for anal fistula” suggested thatthe average rate of incontinence following cutting seton use was 12% while as no incontinence was seen in our study group.

It can be suggested that fistulectomy causes more damage to the sphincter and underlying muscle than fistulotomy. Seton causes minimal damage to the sphincter mechanism and is preferred in high fistulas.

e) Recurrence

Recurrence was noticed in 7% of fistulotomy patients in follow up to 1 year. No similar complaints were noticed in other groups.

Khalid Hussain Qureshi, Mustafa Kamal, Muhammad Hamid Ali Shah, Naseer Ahmed Tariq, Salman Ahmad Tipu in their study “Management of Fistula-in-Ano” had demonstrated overall recurrence rate of 4.44% for low fistulae operated by laying open technique (fistulotomy)and 11.11 % for high fistula in-ano treated with seton cut-through technique. Fistulotomy has higher

incidence of recurrence than fistulectomy as more tissue is left behind. Seton has low rate or recurrence and was nil in the present study.

CONCLUSION

Fistula in ano is an important, commonest disease due to cryptoglandular infection (anal glands) and has a complication of ano rectal abscess.

It is curable disease by the treatment of surgery and higher antibiotics, local antibiotics with good post operative wound management, like sitz bath for twice a day without closing the wound.

Diagnosis is by history, clinical examination, per rectal examination with discharging sinus and pain the complaints in majority of patients.

All the cases should undergo surgery. Fistulotomy although has a slightly higher recurrence rate thanfistulectomy is preferred for low anal fistulas, as it is associated with less chances of incontinence, has significantly less incidence of post operative complications and is associated with less hospital stay duration. Seton happens to be the procedure of choice in high anal fistula.

REFERENCES:

1. Ani A.N. Lagundoye SB, Radiological evaluation of anal fistulae, A prospective study of fistulogram, clinical radiology, 1979 vol. 30 P.21 - 24.
2. Alan. M. Cuthbertson, Sir Edward Hughes, Mark K. Killingback - Anorectal suppuration II - Anal fistula Colorectal surgery, Churchill Livingstone 1983, P - 142 - 162.
3. Buchan, Randgrace R.H. Anorectal suppuration, Results of treatment and factors influencing the recurrence rate British Journal of surgery - 1973 Vol 60, P.537 - 540.
4. Charles Rob and Rodney Smith, clinical surgery Abdomen, Rectum and anus.
5. Chaterjee B.P. Fistula in Ano, A short text book of surgery, Vol 2. II Edn. New Central India Agency.
6. Cyril A. Keele, Eric Neil - Samson Wright, Applied Physiology.
7. Das, S. Clinical Methods in Surgery.
8. Das, S. Operations on rectum and anal canal, A practical guide to operative surgery.
9. Eric L. Farqhharson, R.F. Rintaul, Text book of operative surgery, Churchill Livingstone.
10. Fielding and Garrison, An introducing to history of medicine.
11. Gabriel W.B., The principles and practice of Rectal surgery.
12. Goligher J.C. Surgery of anus, rectum & colon.
13. Harding Rains A.J., Charles V.Mann. Bailey and love's short practice of surgery 23rd Edn.
14. Hartley Stern, Robin Mc Leod, zane Cohen and Theodore Ross, Ano rectal sepsis - Diagnosis and treatment of fistula in ano, Ambulatory procedures in anorectal surgery, advances in surgery, John A. Mannick Editor, Advance in surgery, year book Medical publishers 1987, P.232 - 234.
15. Henry R. Thomson, Henry Souttar, Rectum and anus, British textbook of surgery, Vol.1.
16. James P.S. Thomson, David C. Sabiston Jr. The rectum and anal canal, Text book of surgery.
17. Kron Borg O. - To lay open or to excise a fistula in Ano?, Randomised trial, British Journal of surgery, 1985, Vol. 72. P.970.
18. Lockhart - Mummery H.E. - Charles Rob, Rob smith, Sir, Clifford Naughton Morgan - Operative surgery, Abdomen rectum and anus part II.
19. Parks. A.G. The pathogenesis and treatment of fistula in ano – British medical journal, Vol I 1961, P.463 - 469.
20. Parks. A.G., P.H. Gordon & J.D. hardcastle - A classification of fistula in Ano - British journal of surgery, 1976, Vol.63, P1 - 12.
21. Peter Williams, Roger Warwick, Mary Dyson, Lawrence H. Bannister - Gray's Anatomy 37th Edn. Churchill Livingstone P. 1369 - 1373.
22. Seymour I. Shwartz, Herold Ellis, Rodney Maingot - Abdominal Operation, Vol II.